

## Case Report

## Percutaneous Coil Embolisation of a False Aortic Aneurysm Following Coronary Surgery and Mediastinitis

NIKOLAOS BARBETAKIS<sup>1</sup>, THEOCHARIS XENIKAKIS<sup>2</sup>, ANDREAS EFSTATHIOU<sup>3</sup>, IOANNIS FESSATIDIS<sup>3</sup>

<sup>1</sup>Department of Thoracic Surgery, Theagenio Hospital, Thessaloniki, <sup>2</sup>Department of Cardiothoracic Surgery, University of Crete, Heraklion, <sup>3</sup>Department of Cardiothoracic Surgery, Geniki Kliniki, Thessaloniki, Greece.

Key words:

**Aortic pseudoaneurysm, mediastinal infection.**

A 71-year-old male patient was admitted with methicillin-resistant *staphylococcus aureus* mediastinitis two months after coronary artery bypass grafting. Treatment with immediate surgical debridement, removal of sternal wires and use of vacuum-assisted closure device was started. Spiral computerised tomography and aortography revealed a false aortic aneurysm at the cannulation site. Active mediastinitis and the patient's objection to further surgery led us to perform percutaneous coil embolisation. No postoperative complication was observed and one year later the patient is in excellent condition.

Manuscript received:  
January 1, 2007;  
Accepted:  
March 30, 2007.

Address:

Theocharis Xenikakis

Department of  
Cardiothoracic  
Surgery  
Heraklion University  
Hospital,  
Crete, Greece  
e-mail:  
[txenikakis@yahoo.gr](mailto:txenikakis@yahoo.gr)

**T**hough a rare complication of cardiac surgery, the management of false aortic aneurysms associated with mediastinal infection remains a challenging surgical problem with high associated mortality. Early diagnosis is essential, given the fact that false or mycotic aneurysms progressively expand, compress and erode the surrounding structures and the most significant are associated with a high risk of sudden rupture.

### Case presentation

A 71-year-old male patient with a history of diabetes mellitus type 2, hypertension and chronic obstructive pulmonary disease presented with unstable angina. Coronary angiography revealed three-vessel disease and an ejection fraction of 60%. Subsequently, the patient underwent triple coronary artery bypass grafting. The postoperative period was uneventful and the patient was discharged home on the ninth postoperative day.

On the 60th postoperative day the patient was admitted to the hospital because

of high fever (39.2°C), dyspnoea, chest pain, fatigue, sternal instability and signs of inflammation on the sternotomy wound. Laboratory findings were normal except for an increased leukocyte count (17,000/mm<sup>3</sup>, 88% neutrophils).

The patient underwent urgent surgical debridement under aseptic conditions in the operating theatre. Reopening the wound revealed sternal dehiscence with fragmentation. After removal of the sternal wires, severe suppurative mediastinal infection was confirmed. Probes for bacteriological cultures as well as sternal bone biopsies were taken. Aggressive debridement followed, with removal of all necrotic tissue and irrigation with dilute povidone-iodine solution and H<sub>2</sub>O<sub>2</sub>. Bony debridement was performed until healthy bleeding bone was revealed. Following the debridement procedure, the wound was fitted with a vacuum assisted device.


Computed tomography revealed sternal fragmentation and loss of the integrity of the retrosternal soft tissue fat, indicating mediastinitis as well as a false aortic aneurysm (Figure 1). Aortography confirmed



**Figure 1.** Sternal fragmentation and loss of the integrity of the retrosternal soft tissue fat, indicating mediastinitis as well as a false aortic aneurysm.

the presence of a 20 mm false aneurysm at the cannulation site (Figure 2).

Advanced age, active mediastinitis and the patient's objection to further surgery led us to propose a palliative percutaneous treatment. Via femoral access, through a 15 cm long 8F diameter sheath and a 7.2F diameter guiding catheter, a balloon catheter was placed so as to occlude the false aneurysm neck on balloon inflation (Moret's neurointerventional technique).<sup>1</sup> Three long coils of spiroid shape were mechanically dropped through the catheter lumen. Stable homogeneous packing was achieved and the balloon was progressively deflated after testing the material's stability



**Figure 2.** Aortography confirmed the presence of a 20 mm false aneurysm at the cannulation site.

ty by successful injection into the ascending aorta (Figure 3).

The patient did well with no postoperative complications. Serial quantitative wound cultures were positive for methicillin-resistant *staphylococcus aureus* and intravenous antibiotic therapy was started (vancomycin 500 mg × 3 daily). The vacuum assisted closure system was changed every two to three days and used in total for 22 days. The system was removed when regional and systemic signs of infection resolved and quantitative wound cultures were negative. Finally, the patient underwent regional muscle flap closure (pectoralis flap) and complete healing was achieved. One year later he is in excellent condition and magnetic resonance imaging confirmed coil stability.

### Discussion

Infectious mediastinitis is a relatively common but potentially devastating complication occurring after cardiac operations that are performed through median sternotomy. Despite an incidence of less than 5%, the importance of this complication should not be underestimated.<sup>2</sup> Infectious mediastinitis with pseudoaneurysm formation is a very rare but highly morbid complication due to subsequent rupture.<sup>3</sup> Without an operation, aortic false aneurysms progressively expand, compress and erode the surrounding structures, or are a source of persistent infection and systemic embolism. Percutaneous embolisation of large saccular aneurysms or pseudoaneurysms is also effective.<sup>4,5</sup>



**Figure 3.** Stable homogeneous packing of the aneurysm after occlusion by coils.

In the case reported here, active mediastinitis and the patient's reluctance to undergo another operation led us to perform percutaneous treatment according to the Moret remodelling technique. This technique consists of inflating a balloon in front of the neck, thus allowing the coils to be pushed into the aneurysm while avoiding any outside bulging. The Moret technique has been used successfully to treat 56 intracranial wide-necked or badly-shaped aneurysms with satisfactory occluder stability during follow up.<sup>1</sup>

In conclusion, the treatment of false aneurysms poses special challenges to the cardiothoracic surgeon. The optimal surgical management is a subject of controversy. Endovascular treatment with coils may be a good and safe alternative to surgery.

## References

1. Moret J, Cognard C, Weill A, Castaing L, Rey A: The remodeling technique in the treatment of wide neck intracranial aneurysms: angiographic and clinical follow up about 56 cases. *Intervent Neuroradiol* 1997; 3: 21-25.
2. King RC, Barnes AD: Mediastinitis after cardiac surgery. *Curr Treat Options Infect Dis* 2003; 5: 377-386.
3. Vrochides D, Feng WC, Singh AK: Mycotic ascending aortic pseudoaneurysm secondary to pseudomonas mediastinitis at the aortic cannulation site. *Tex Heart Inst J* 2003; 30: 322-324.
4. Rao VR, Mandalam RK, Joseph S, et al: Embolization of large saccular aneurysms with Gianturco coils. *Radiology* 1990; 175: 407-410.
5. Miguel B, Camilleri L, Gabrillargues J, et al: Coil embolization of a false aneurysm with aortocutaneous fistula after prosthetic graft replacement of the ascending aorta. *Eur J Radiol* 2000; 34: 57-59.